AIDS Surveillance: Demographics of All Diagnosed Cases - 2010

Cumulative^ Percent of Reported AIDS Cases Erie County, PA, and U.S.

		ERIE COUNTY	PA	U.S.	U.S.
		% of Total Reported	% of Total Reported	% of Total Reported	% of Total Estimated**
		As of 12/31/2010	As of 12/31/2010	As of 12/31/2009	As of 12/31/2009
Gender	Male*	82.8	77.5	79.9	79.8
	Female*	17.2	22.5	20.1	20.2
Age at Diagnosis	<13 (Pediatric)	0.8	1.0	0.9	0.9
	13-19	0.8	0.9	0.8	0.8
	20-29	19.2	15.1	15.6	15.6
	30-39	40.3	38.6	40.7	40.5
	40-49	27.6	29.9	28.7	28.8
	≥ 50	11.3	14.6	13.4	13.5
Race / Ethnicity	White, non-Hispanic	58.6	35.5	37.6	37.3
	Black, non-Hispanic	29.9	49.6	40.7	40.9
	Hispanic	11.3	12.4	19.5	19.6
	Other	0.3	2.5	2.2	2.2
	Unknown	0.0	0.0	0.0	0.0
Mode of Transmission	Men who have sex with men (MSM)	43.5	37.6	44.9	47.6
	Injecting drug use (IDU)	25.1	30.5	23.2	25.7
	MSM & IDU	6.9	5.1	6.8	7.1
	High risk heterosexual contact	15.3	20.5	15.1	18.5
	Hemophilia/Coagulation disorder	0.9	0.7	n/a	n/a
	Transfusion/Receipt of blood components or tissue	1.2	0.6	n/a	n/a
	Other/ Risk not reported***	6.1	1.1	9.1	1.1
	All Pediatric****	0.0	0.0	0.8	0.8

[^]From beginning of epidemic (1980 for PA and Erie; 1983 for U.S.)

Data may be incomplete due to a reporting lag; Number of cases listed in this report may vary from previous reports due to the interstate deduplication process United States cases represent 50 states, District of Columbia, and U.S. dependent areas

Sources: Erie County Department of Health; PA DOH; PA DOH HIV Surveillance Summary Report, 2010; CDC HIV/AIDS Surveillance Reports, 2009



^{*}For PA, gender is reported for all age groups; For U.S., only gender for adults and adolescents is reported

^{**}Estimating cases accounts for reporting delays

^{***}For U.S., includes hemophilia, blood transfusion, and any other risk factor not reported or not identified

^{****}Includes adult cases which are assigned pediatric modes of transmission since infection is believed to have occurred before age 13 n/a=Not available